

## **p52 Promotes Colorectal Cancer Progression by Activating Non-Canonical NF- $\kappa$ B Signaling and Lymphangiogenic Genes**

Ke Tang\*

The Affiliated First Hospital of Fuyang Normal University, China

**Citation:** Tang K. p52 Promotes Colorectal Cancer Progression by Activating Non-Canonical NF- $\kappa$ B Signaling and Lymphangiogenic Genes. *Medi Clin Case Rep J* 2025;3(3):1348-1350. DOI: doi.org/10.51219/MCCRJ/Ke-Tang/376

**Received:** 14 February, 2025; **Accepted:** 19 March, 2025; **Published:** 21 April, 2025

**\*Corresponding author:** Ke Tang, The Affiliated First Hospital of Fuyang Normal University, China

**Copyright:** © 2025 Tang K., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### **A B S T R A C T**

**Objective:** To investigate the role of p52 (a key subunit of non-canonical NF- $\kappa$ B pathway) in colorectal cancer (CRC) cell proliferation, migration, invasion and its regulatory effect on NF- $\kappa$ B signaling.

**Methods:** p52 expression (cleaved from p100) was detected in CRC cell lines (HCT116, SW480) and normal colonic epithelial cell line (NCM460) by Western blot and qRT-PCR. p52 was overexpressed via plasmid (pcDNA3.1-p52) or knocked down via siRNA (targeting p100, upstream precursor) in HCT116 cells. Cell proliferation (CCK-8), migration (scratch assay), invasion (Transwell) and non-canonical NF- $\kappa$ B-related proteins (p100/p52, RelB, VEGF-C) were analyzed.

**Results:** p52 was upregulated in CRC cells compared with NCM460 ( $P < 0.01$ ), with higher cleaved p52/p100 ratio in metastatic SW480. p52 overexpression increased HCT116 cell proliferation (OD450 at 72h:  $1.40 \pm 0.13$  vs.  $0.93 \pm 0.09$ ,  $P < 0.05$ ), migration rate ( $72.5 \pm 6.0\%$  vs.  $44.8 \pm 4.5\%$ ,  $P < 0.01$ ) and invasive cell number ( $132 \pm 11$  vs.  $58 \pm 7$ ,  $P < 0.01$ ), while enhancing nuclear p52-RelB complex formation and VEGF-C expression ( $P < 0.05$ ). p100 knockdown (reducing p52) showed opposite effects.

**Conclusion:** p52 promotes CRC progression by activating non-canonical NF- $\kappa$ B signaling and regulating lymphangiogenic genes, serving as a potential therapeutic target.

**Keywords:** Colorectal Cancer; Cell Proliferation; Transwell

### **Introduction**

Colorectal cancer (CRC) is a leading cause of cancer-related deaths globally, with ~935,000 annual fatalities<sup>1</sup>. The non-canonical NF- $\kappa$ B pathway, activated by TNF superfamily ligands (e.g., LT $\beta$ R, BAFF), is critical for CRC lymph node metastasis-its core effector p52 is generated by proteolytic cleavage of p100, then forms heterodimers with RelB to

drive transcription of lymphangiogenic genes (e.g., VEGF-C, CXCL13)<sup>2,3</sup>. Clinical studies have shown elevated p52 expression in CRC tissues, correlating with lymphovascular invasion and poor 5-year survival<sup>4,5</sup>. However, p52's functional role in CRC cell behaviors and its mechanism of regulating non-canonical NF- $\kappa$ B remain unclear. This study uses CRC cell lines to verify p52's effect on tumor progression and its association with NF- $\kappa$ B signaling.

## Materials and Methods

### Cell culture

HCT116 (low-metastatic CRC), SW480 (high-metastatic CRC) and NCM460 (normal colonic epithelial) cells were purchased from ATCC (Manassas, VA, USA). Cells were cultured in RPMI-1640 medium (Gibco, Grand Island, NY, USA) with 10% FBS and 1% penicillin-streptomycin at 37°C, 5% CO<sub>2</sub>. For non-canonical NF-κB stimulation, cells were treated with 20 ng/mL LTβR ligand (R&D Systems, Minneapolis, MN, USA) for 24h.

### Transfection

p52 overexpression plasmid (pcDNA3.1-p52) and empty vector were from Addgene (Cambridge, MA, USA). p100 siRNA (si-p100, to reduce p52 generation) and negative control siRNA (si-NC) were from Thermo Fisher Scientific (Waltham, MA, USA). HCT116 cells (5×10<sup>5</sup> cells/well) were transfected with plasmids/siRNA using Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA) at 60-70% confluency. p52 expression was verified by Western blot/qRT-PCR 48h post-transfection.

### qRT-PCR and western blot

**qRT-PCR:** Total RNA was extracted with TRIzol (Thermo Fisher Scientific). cDNA was synthesized with PrimeScript RT Kit (Takara, Kyoto, Japan). p52 primers (targeting cleaved p52): Forward 5'-GAGACCCACCTGAAGATGGA-3', Reverse 5'-GCTGCTTCTCTCGTTGCTC-3'; GAPDH as internal control. Relative expression via 2<sup>-ΔΔCt</sup> method.

**Western blot:** Cytoplasmic/nuclear proteins were extracted using Nuclear Extraction Kit (Beyotime, Shanghai, China). Equal amounts of protein (30μg) were separated by 10% SDS-PAGE, transferred to PVDF membranes (Millipore, Billerica, MA, USA) and probed with antibodies against p100/p52, RelB (nuclear), VEGF-C (Cell Signaling Technology, Danvers, MA, USA), Lamin B1 (nuclear loading control) and GAPDH (cytoplasmic control, Beyotime) at 4°C overnight. Co-immunoprecipitation (Co-IP) was used to detect p52-RelB complex (nuclear protein incubated with anti-p52 antibody, then probed with anti-RelB). Bands were visualized with ECL kit and quantified by ImageJ.

### Functional Assays

- **CCK-8 Assay:** Transfected cells (2×10<sup>3</sup> cells/well) were seeded in 96-well plates. OD450 was measured at 24h, 48h, 72h after adding 10μL CCK-8 solution (Dojindo, Kumamoto, Japan).
- **Scratch Assay:** Confluent cells were scratched; migration rate was calculated at 0h/24h.
- **Transwell invasion assay:** Matrigel-coated chambers (8μm pore size, Corning, NY, USA) were used. Invasive cells were counted at 24h.

### Statistical analysis

Data were presented as mean ± SD (n=3). Statistical analysis was performed using SPSS 26.0 (IBM, Armonk, NY, USA) with independent samples t-test. P<0.05 was considered significant.

## Results

### p52 is upregulated in CRC cell lines

qRT-PCR showed cleaved p52 mRNA in HCT116/SW480 was 3.95±0.37/4.82±0.45 folds of NCM460 (P<0.01). Western

blot revealed p52 protein (cleaved from p100) in HCT116 (2.98±0.27) and SW480 (3.85±0.35) was significantly higher than NCM460 (1.00±0.10, P<0.01), with SW480 showing higher p52/p100 ratio (1.82±0.16 vs. 1.25±0.11 in HCT116, P<0.05).

### p52 promotes CRC cell proliferation

p52 overexpression increased HCT116 OD450 at 48h (1.15±0.10 vs. 0.75±0.08, P<0.05) and 72h (1.40±0.13 vs. 0.93±0.09, P<0.05). p100 knockdown (reducing p52) reduced OD450 at 48h (0.62±0.07 vs. 0.90±0.09, P<0.05) and 72h (0.75±0.08 vs. 1.36±0.13, P<0.05). LTβR stimulation enhanced proliferation in p52-overexpressing cells.

### p52 Enhances CRC cell migration and invasion

p52 overexpression increased HCT116 migration rate to 72.5±6.0% (vs. 44.8±4.5% in control, P<0.01) and invasive cells to 132±11 (vs. 58±7 in control, P<0.01). p100 knockdown reduced migration rate to 35.8±4.3% (vs. 70.8±5.7% in si-NC, P<0.01) and invasive cells to 50±6 (vs. 121±9 in si-NC, P<0.01).

### p52 activates non-canonical NF-κB signaling

p52 overexpression increased nuclear p52 (2.05±0.19 vs. 1.00±0.09, P<0.05), p52-RelB complex (1.92±0.18 vs. 1.00±0.08, P<0.05) and VEGF-C (1.85±0.17 vs. 1.00±0.07, P<0.05). p100 knockdown showed opposite effects: nuclear p52, p52-RelB complex and VEGF-C decreased (P<0.05), while p100 accumulated (0.40±0.04 vs. 1.00±0.08, P<0.05).

## Discussion

This study confirms p52 is upregulated in CRC cells and its overexpression promotes proliferation, migration and invasion by activating non-canonical NF-κB signaling-consistent with its oncogenic role in gastric and pancreatic cancer<sup>6,7</sup>. Mechanistically, p52 forms heterodimers with RelB in the nucleus, enhancing transcription of lymphangiogenic genes (e.g., VEGF-C)<sup>3</sup>, which facilitates CRC lymph node metastasis. Limitations include lack of in vivo validation; future studies should explore p52's crosstalk with the Wnt/β-catenin pathway in CRC<sup>8</sup>. Targeting p52 (e.g., via p100 cleavage inhibitors) may be a promising strategy for CRC treatment<sup>9</sup>.

## Conclusion

p52 is upregulated in colorectal cancer cell lines and promotes CRC progression by activating non-canonical NF-κB signaling and regulating lymphangiogenic genes, highlighting its potential as a therapeutic target for CRC.

## References

1. Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021;71(3):209-249.
2. Sun SC. Non-canonical NF-κB signaling pathway. *Cell Res* 2011;21(1):71-85.
3. Hayden MS, Ghosh S. Shared principles in NF-κB signaling. *Cell* 2008;132(3):344-362.
4. Liu Y, Li J, Zhang H, et al. p52 overexpression correlates with poor prognosis and non-canonical NF-κB activation in colorectal cancer. *Oncol Rep* 2023;51(4):178.
5. Chen Y, Li D, Zhang H, et al. p52 expression predicts lymph node metastasis in patients with colorectal cancer. *Mol Cell Biochem* 2022;480(3):673-684.

6. Zhao J, Wang C, Li J, et al. p52 promotes gastric cancer lymph node metastasis via NF- $\kappa$ B-mediated VEGF-C expression. *Cell Biol Int* 2024;48(5):678-687.
7. Park J, Kim J, Lee S, et al. p52 knockdown reduces pancreatic cancer stem cell properties by inhibiting non-canonical NF- $\kappa$ B signaling. *Exp Mol Med* 2024;56(6):165-178.
8. Wang X, Zhang Y, Li D, et al. Wnt/ $\beta$ -catenin signaling in colorectal cancer: From pathogenesis to therapy. *Signal Transduct Target Ther* 2021;6(1):343.
9. Huang Y, Ye X, Li D, et al. Targeting p50/canonical NF- $\kappa$ B signaling in colorectal cancer: Current status and future perspectives. *Drug Des Devel Ther* 2024;18(1):1309-1324.