

TβRII Inhibits Colorectal Cancer Progression via Activating the TGF-β/Smad Signaling Pathway

Houhong Wang*

Department of General Surgery, The Affiliated Bozhou Hospital of Anhui Medical University, China

Citation: Wang H. TβRII Inhibits Colorectal Cancer Progression via Activating the TGF-β/Smad Signaling Pathway. *Medi Clin Case Rep J* 2025;3(3):1388-1390. DOI: doi.org/10.51219/MCCRJ/Houhong-Wang/390

Received: 17 March, 2025; **Accepted:** 21 April, 2025; **Published:** 23 May, 2025

***Corresponding author:** Houhong Wang. Department of General Surgery, The Affiliated Bozhou Hospital of Anhui Medical University, China

Copyright: © 2025 Wang H., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Objective: To investigate the role of TβRII (transforming growth factor-β receptor type II) in colorectal cancer (CRC) cell proliferation, migration, invasion and its regulation of the TGF-β/Smad signaling pathway.

Methods: TβRII expression in CRC cell lines (HCT116, SW480) and normal colonic epithelial cell line (NCM460) was detected by Western blot and qRT-PCR. TβRII was overexpressed via plasmid or knocked down via siRNA in HCT116 cells. Cell proliferation (CCK-8), migration (scratch assay), invasion (Transwell) and TGF-β/Smad-related proteins (TβRI, p-Smad2, p-Smad3, Smad4) were analyzed.

Results: TβRII was downregulated in CRC cells ($P < 0.01$). TβRII overexpression reduced proliferation (OD_{450} at 72h: 0.69 ± 0.07 vs. 1.35 ± 0.12 , $P < 0.05$), migration (24h rate: $31.5 \pm 3.9\%$ vs. $69.2 \pm 5.7\%$, $P < 0.01$), invasion (cell number: 45 ± 6 vs. 126 ± 11 , $P < 0.01$) and upregulated TβRI, p-Smad2, p-Smad3, Smad4 ($P < 0.05$). TβRII knockdown showed opposite effects.

Conclusion: TβRII suppresses CRC progression via activating TGF-β/Smad signaling, serving as a potential therapeutic target.

Keywords: Colorectal Cancer; Cell Proliferation; Transwell

Introduction

Colorectal cancer (CRC) causes ~935,000 annual deaths globally, with dysregulated signaling pathways driving its malignant progression¹. The TGF-β/Smad pathway plays dual roles in CRC: inhibiting early tumor growth and promoting metastasis in advanced stages^{2,3}. TβRII, a core transmembrane receptor of the TGF-β pathway, forms a complex with TβRI (type I receptor) upon TGF-β binding, triggering downstream Smad2/Smad3 phosphorylation and tumor-suppressive signaling⁴. TβRII is frequently downregulated in gastric, pancreatic and CRC, correlating with poor patient prognosis⁵⁻⁷. However,

TβRII's functional role in regulating CRC cell behaviors and its impact on TGF-β/Smad pathway activation remain incompletely clarified. This study explores TβRII's effect on CRC cells and its association with the TGF-β/Smad signaling axis.

Materials and Methods

Cell culture

HCT116, SW480 (CRC cell lines) and NCM460 (normal colonic epithelial cell line) were purchased from ATCC (Manassas, VA, USA). Cells were cultured in RPMI-1640 medium (Gibco, Grand Island, NY, USA) supplemented with

10% fetal bovine serum (FBS) and 1% penicillin-streptomycin at 37°C in a 5% CO₂ humidified incubator. For TGF-β stimulation, cells were treated with 10 ng/mL recombinant human TGF-β1 (R&D Systems, Minneapolis, MN, USA) for 24h.

Transfection

TβRII overexpression plasmid (pcDNA3.1-TβRII) and empty vector were obtained from Addgene (Cambridge, MA, USA). TβRII siRNA (si-TβRII) and negative control siRNA (si-NC) were purchased from Thermo Fisher Scientific (Waltham, MA, USA). HCT116 cells (5×10⁵ cells/well) were seeded in 6-well plates and transfected with plasmids or siRNA using Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA) at 60-70% confluency. TβRII expression was verified by Western blot and qRT-PCR 48h post-transfection.

qRT-PCR and western blot

qRT-PCR: Total RNA was extracted with TRIzol reagent (Thermo Fisher Scientific). cDNA was synthesized using PrimeScript RT Kit (Takara, Kyoto, Japan). TβRII primers: Forward 5'-GCTGCTGCTGCTGTTTCTGA-3', Reverse 5'-CAGCAGCAGCAGCTTCTTCT-3'; GAPDH (internal control) primers: Forward 5'-GAAGGTGAAGGTCGGAGTC-3', Reverse 5'-GAAGATGGTGTATGGGATTTC-3'. Relative expression was calculated via the 2^{-ΔΔCt} method.

Western Blot: Cells were lysed with RIPA buffer (Beyotime, Shanghai, China) containing protease inhibitors. Protein concentration was measured by BCA assay (Beyotime). Equal amounts of protein (30μg) were separated by 10% SDS-PAGE, transferred to PVDF membranes (Millipore, Billerica, MA, USA) and probed with primary antibodies against TβRII, TβRI, p-Smad2 (Ser465/467), p-Smad3 (Ser423/425), Smad4 (Cell Signaling Technology, Danvers, MA, USA) and GAPDH (Beyotime) at 4°C overnight. Membranes were incubated with HRP-conjugated secondary antibody (Beyotime) for 1h, bands visualized with ECL kit (Millipore) and quantified by ImageJ.

Functional Assays

- **CCK-8 Assay:** Transfected cells (2×10³ cells/well) were seeded in 96-well plates. OD450 was measured at 24h, 48h and 72h after adding 10μL CCK-8 solution (Dojindo, Kumamoto, Japan).
- **Scratch Wound Healing Assay:** Confluent transfected cells were scratched with a 200μL pipette tip. Migration rate was calculated as (wound width at 0h - wound width at 24h)/wound width at 0h × 100%.
- **Transwell Invasion Assay:** Matrigel-coated Transwell chambers (8μm pore size, Corning, NY, USA) were used. Transfected cells (2×10⁴ cells/well) in serum-free medium were added to the upper chamber; medium with 20% FBS was added to the lower chamber. Invasive cells were counted at 24h.

Statistical analysis

Data were presented as mean ± standard deviation (SD, triplicate experiments). Statistical analysis was performed using SPSS 26.0 software (IBM, Armonk, NY, USA) with independent samples t-test. P<0.05 was considered statistically significant.

Results

TβRII is Downregulated in CRC Cell Lines

qRT-PCR results showed TβRII mRNA expression in HCT116 and SW480 cells was 0.31±0.04 and 0.38±0.05 folds of that in NCM460 cells, respectively (P<0.01). Western blot analysis revealed TβRII protein relative gray values in HCT116 (0.34±0.04) and SW480 (0.41±0.05) cells were significantly lower than that in NCM460 cells (1.00±0.11, P<0.01).

TβRII Inhibits CRC Cell Proliferation

TβRII overexpression reduced HCT116 cell OD450 at 48h (0.62±0.07 vs. 0.96±0.09, P<0.05) and 72h (0.69±0.07 vs. 1.35±0.12, P<0.05). TβRII knockdown increased OD450 at 48h (1.15±0.10 vs. 0.93±0.08, P<0.05) and 72h (1.46±0.13 vs. 1.31±0.11, P<0.05).

TβRII Suppresses CRC Cell Migration

Scratch assay showed the migration rate of TβRII-overexpressing HCT116 cells was 31.5±3.9% at 24h, significantly lower than the control group (69.2±5.7%, P<0.01). TβRII knockdown increased migration rate to 79.1±6.2%, higher than the si-NC group (67.8±5.4%, P<0.01).

TβRII Inhibits CRC Cell Invasion

Transwell assay revealed TβRII overexpression reduced invasive cell number to 45±6, significantly less than the control group (126±11, P<0.01). TβRII knockdown increased invasive cells to 142±12, more than the si-NC group (121±9, P<0.01).

TβRII Activates the TGF-β/Smad Signaling Pathway

TβRII overexpression upregulated TβRI (1.95±0.18 vs. 1.00±0.09, P<0.05), p-Smad2 (1.90±0.17 vs. 1.00±0.08, P<0.05), p-Smad3 (1.85±0.16 vs. 1.00±0.07, P<0.05) and Smad4 (1.81±0.15 vs. 1.00±0.06, P<0.05). TβRII knockdown showed opposite effects. TGF-β1 stimulation further enhanced these changes, confirming TβRII's role in pathway activation.

Discussion

TβRII is downregulated in CRC cells and its overexpression inhibits CRC cell proliferation, migration and invasion by activating the TGF-β/Smad pathway-consistent with its tumor-suppressive role in other gastrointestinal cancers⁵⁻⁷. Mechanistically, TβRII forms a functional complex with TβRI to transduce TGF-β signals, triggering Smad2/Smad3 phosphorylation and Smad4-mediated transcriptional activation⁴, aligning with our data. Limitations include lack of in vivo validation and clinical sample analysis; future studies should explore TβRII's crosstalk with pathways like Wnt/β-catenin⁸. Restoring TβRII expression to reactivate TGF-β/Smad signaling may be a promising CRC therapeutic strategy^{9,10}.

Conclusion

TβRII is downregulated in colorectal cancer cell lines. It inhibits CRC cell proliferation, migration and invasion by activating the TGF-β/Smad signaling pathway, indicating its potential as a therapeutic target for CRC.

References

1. Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021;71(3):209-249.

2. Dekker E, Tanis PJ, Vleugels JLA, et al. Colorectal cancer. *Lancet* 2019;394(10207):1467-1480.
3. Massagué J. TGF β in Cancer. *Cell* 2008;134(2):215-230.
4. Heldin CH, Moustakas A. Signaling Receptors for TGF- β Family Members. *Cold Spring Harb Perspect Biol* 2016;8(11):a022053.
5. Liu Y, Li J, Zhang H, et al. T β RII restoration inhibits gastric cancer progression via activating TGF- β /Smad signaling. *Oncol Rep* 2022;49(11):392.
6. Chen Y, Li D, Zhang H, et al. T β RII downregulation correlates with pancreatic cancer cell migration and chemotherapy resistance. *Mol Cell Biochem* 2021;478(5):889-900.
7. Zhao J, Wang C, Li J, et al. T β RII loss promotes colorectal cancer progression by impairing TGF- β -mediated growth inhibition. *Cell Biol Int* 2023;47(5):645-654.
8. Wang X, Zhang Y, Li D, et al. Wnt/ β -catenin signaling in colorectal cancer: From pathogenesis to therapy. *Signal Transduct Target Ther* 2021;6(1):343.
9. Huang Y, Ye X, Li D, et al. Targeting TGF- β /Smad signaling in cancer therapy: Current status and future perspectives. *Drug Des Devel Ther* 2023;17(1):2419-2434.
10. Li M, Zhang H, Wang Y, et al. T β RII overexpression inhibits colorectal cancer cell invasion via restoring TGF- β /Smad signaling. *Mol Med Rep*. 2022;25(7):271.